

Find a coordinate independent expression for the distance between two points and then evaluate it in rectangular coordinates.

Solution If you are trying to find the distance between two objects, ideally you just measure with a tape measure or equivalent. But sometimes you don't have access to the two objects e.g. they are in outer space or tiny under a microscope). In particular, if you are trying to *calculate* the distance in an equation, then you need to do something more. First you must locate the two objects using their positions vectors \vec{r} and \vec{r}' relative to a common origin. Then you must write the position vectors in a common coordinate system $\vec{r} = x\hat{x} + y\hat{y}$, etc. When you subtract the two vectors (use *vector* subtraction!!) you get the vector that goes from one object to the other, $\vec{r} - \vec{r}' = (x - x')\hat{x} + (y - y')\hat{y}$. You can find the length of this (or any other vector) by finding the square root of the dot product of the vector with itself.

$$|\vec{r} - \vec{r}'| = \sqrt{(\vec{r} - \vec{r}') \cdot (\vec{r} - \vec{r}')} \quad (1)$$

$$= \sqrt{((x - x')\hat{x} + (y - y')\hat{y}) \cdot ((x - x')\hat{x} + (y - y')\hat{y})} \quad (2)$$

$$= \sqrt{(x - x')^2 + (y - y')^2} \quad (3)$$

N.B. Of course, this last expression is the same as the Pythagorean theorem on a triangle whose hypotenuse is the line segment between the two objects. This is a method of PROVING the Pythagorean theorem. It will help you in future problems to know how to do this calculation the long way, described here.